Me know the facts

What are the risks of marijuana use?

Anxiety	Depression

- _ack of motivation
- 0 0 Mood swings
- Schizophrenia
- 0 0 Damages airways
- 0 Asthma Lung infections
- 0 **Emphysema**
- 0 Lung cancer
- 0 Impairs immune system

0 Delayed reaction time and coordination resulting in injury or car crashes

0

Addiction

- 0 Impairs short term memory, learning, judgement, and sensory perception
- 0 pressure, heart rate and risk of heart attack
- Weight gain

MEN: Lowers testosterone and sperm production, increases risk for erectile dysfuntion, breast growth, and testicular cancer

WOMEN: Fertility problems

PREGNANCY: Increased risk of premature birth, THC transfers to baby through placenta and breast milk

Need help?

Washington Recovery Help Line For free referrals and emotional support:

1-866-789-1511

Teen Link

1-866-833-6546

Quick facts

- O **Just like alcohol**, marijuana has health and safety risks.
- O **Just like alcohol**, marijuana can damage the developing brain. Avoid using if pregnant, nursing or under 21.





Increased risk of heart attack in the first hour after smoking



Increased risk for psychosis



Get more facts at: www.LearnAboutMarijuanaWA.org







) know the facts

What are the risks of drinking too much?

- 0 0 Cancers of the throat or mouth Premature aging 0000 00 Depression Anxiety Aggressive behavior Memory loss Insomnia Alcohol dependence
- 000 High blood pressure
- 0 Heart failure

 Blood clotting/low iron levels Breast cancer
- 0 0 00 0 pancreas damage Liver and Malnutrition Bleeding ulcers Stomach inflammation Diarrhea

WOMEN: Deformed, developmentally delayed or low-birth-weight babies

MEN: Erectile dysfunction

0

Increased risk of pneumonia and colds

0 Failure to fulfill obligations at work, school, or home

0

Impaired sensation leading to falls

00 Legal problems Car crashes

Need help?

Washington Recovery Help Line For free referrals and emotional support:

1-866-789-1511

Teen Link 1-866-833-6546

What is a standard drink?

00

Painful nerves
Numb, tingling toes



Low-risk drinking guidelines

All 66+	Women	Men
W	3	Per Day
7	7	Per Week

- If you drink more than this, ask your doctor about ways to cut down.
- Avoid alcohol if pregnant or under 21.



S know the fa

prescription pain relievers (opioids)? What are the risks of misusing

Talk with your health care provider about all the prescription drugs you take.

- Mental confusion
- 000 Drowsiness
- Changes in brain that build tolerance, leading to dependence

WITHDRAWAL SYMPTOMS: Restlessness, muscle and bone pain, insomnia, diarrhea, vomiting, cold flashes with goose bumps, kicking movements, severe craving that can lead to relapse

0

Increased and/or decreased blood pressure

linfectious diseases such as HIV, Hepatitis B and Hepatitis C if injected

TOTAL BODY EFFECTS: Euphoria, warm flushing, poor nutrition, fatal overdose

PREGNANCY AND INFANTS: Miscarriage, low birth weight, developmental delays, addiction

0 HIV and other sexually transmitted infections from unsafe sex while intoxicated 0 0 Nausea Slow, irregular breathing that can be fatal Cramping Constipation

Symptoms of overdose

COLD/CLAMMY SKIN DRY MOUTH WEAK PULSE DROWSINESS LOW BLOOD PRESSURE COMA BLUE NAILS AND LIPS MUSCLE SPASMS DISORIENTATION VOMITING NON RESPONSIVE SEIZURES SMALL PUPILS NO BREATHING OR SHALLOW BREATHING

Need help?

For free referrals and emotional support:

1-866-833-6546

Washington Recovery Help Line 1-866-789-1511 Teen Link

Quick facts



- Prescription opioids are pain. medications that relieve
- 0 They can be addictive if your doctor. not used as directed by
- 0 More people die from opioid overdoses than from any other drug.



Nearly 17,000 people die each year from prescription pain reliever overdoses.

Lock up your meds and safely discard them. Visit: www.TakeBackYourMeds.org



