

prescription pain relievers (opioids)? What are the risks of misusing

Talk with your health care provider about all the prescription drugs you take.

- Mental confusion
- 000 **Drowsiness**
- Changes in brain that build tolerance, leading to dependence

linfectious diseases such as HIV, Hepatitis B and Hepatitis C if injected

0

Restlessness, muscle and bone pain, insomnia, diarrhea, vomiting, cold flashes with goose bumps, kicking movements, severe craving that WITHDRAWAL SYMPTOMS:

TOTAL BODY EFFECTS: Euphoria, warm flushing, poor nutrition, fatal overdose

PREGNANCY AND INFANTS: Miscarriage, low birth weight, developmental delays, addiction

0 HIV and other sexually transmitted infections from unsafe sex while intoxicated 0 0 Nausea Cramping Constipation Slow, irregular breathing that can be fatal Increased and/or decreased blood pressure

Symptoms of overdose

WEAK PULSE DROWSINESS LOW BLOOD PRESSURE COMA BLUE NAILS AND LIPS MUSCLE SPASMS DISORIENTATION VOMITING NON RESPONSIVE SEIZURES SMALL PUPILS NO BREATHING
SHALLOW BREATHING COLD/CLAMMY SKIN DRY MOUTH

Need help?

Washington Recovery Help Line For free referrals and emotional support:

1-866-789-1511

1-866-833-6546 Teen Link

Quick facts



- O Prescription opioids are pain. medications that relieve
- 0 They can be addictive if your doctor. not used as directed by
- 0 More people die from opioid overdoses than from any other drug.



Nearly 17,000 people die each year from prescription pain reliever overdoses.

Lock up your meds and safely discard them. Visit: www.TakeBackYourMeds.org



